

THE HODGE RANK OF ACM BUNDLES AND FRANCHETTA'S CONJECTURE

INDRANIL BISWAS AND G. V. RAVINDRA

ABSTRACT. We prove that on a general hypersurface in \mathbb{P}^N of degree d and dimension at least 2, if an arithmetically Cohen-Macaulay (ACM) bundle E and its dual have small regularity, then any non-trivial Hodge class in $H^n(X, E \otimes \Omega_X^n)$, $n = \lfloor \frac{\dim X}{2} \rfloor$, produces a trivial direct summand of E . As a consequence, we prove that there is no universal Ulrich bundle on the family of smooth hypersurfaces of degree $d \geq 3$ and dimension at least 4. This last statement may be viewed as a Franchetta-type conjecture for Ulrich bundles on smooth hypersurfaces.

1. INTRODUCTION

The Franchetta conjecture, first stated in [13] and proved in [16] (see also [1]), says that for the universal family of genus $g \geq 2$ curves

$$\pi : \mathcal{C}_g \longrightarrow \mathcal{M}_g,$$

the restriction of any line bundle \mathcal{L} on \mathcal{C}_g to any smooth fiber $C_s := \pi^{-1}(s)$ is a power of the canonical bundle.

Since then, various analogues of Franchetta's conjectures have been posed. For instance, motivated by the work of Beauville and Voisin on the Chow groups of $K3$ surfaces (see [8]), O'Grady [26] conjectured that for a smooth family of $K3$ surfaces $\mathcal{X} \rightarrow S$, the restriction map of rational Chow groups $\mathrm{CH}^2(\mathcal{X}) \rightarrow \mathrm{CH}^2(X)$, where X is any smooth fiber, is generated by the class \mathfrak{o}_X – this is the class of a point lying on a rational curve C in X . There are other such generalizations to Chow groups of higher codimensional cycles for the hyper-Kähler varieties (see [9]).

In this article, we study a Franchetta type conjecture in a different direction, *viz. for higher rank bundles* on smooth projective varieties.

Recall that a vector bundle E on a polarized projective variety $(X, \mathcal{O}_X(1))$ is said to be *arithmetically Cohen-Macaulay* (ACM) if it has no intermediate cohomology, meaning

$$H_*^i(X, E) := \bigoplus_{\nu \in \mathbb{Z}} H^i(X, E(\nu)) = 0 \quad \forall 0 < i < \dim X.$$

A result of Horrocks [18] states that a vector bundle on \mathbb{P}^N is a direct sum of line bundles if and only if it is ACM. This statement however does not hold for other classes of varieties in this generality (see for example Proposition 5 in [25]). More recently, a special class of ACM bundles, namely that of *Ulrich bundles*, has received a considerable amount of interest. One striking result that has been proved is that every smooth complete

2010 *Mathematics Subject Classification.* 14J60, 14D21.

Key words and phrases. Franchetta conjecture, ACM bundle, Ulrich bundle, Noether-Lefschetz theorem, Hodge cohomology.

intersection variety over an algebraically closed field supports an Ulrich bundle (see [4]). We recall the following characterization of Ulrich bundles.

Proposition 1.1 (see Theorem 2.3, [5]). *Let $X \subset \mathbb{P}^N$ be a smooth projective variety, and let E be a vector bundle on X . The following conditions are equivalent:*

(i) *There exists a linear resolution*

$$0 \longrightarrow \tilde{L}_c \longrightarrow \tilde{L}_{c-1} \longrightarrow \cdots \longrightarrow \tilde{L}_0 \longrightarrow E \longrightarrow 0,$$

where $c := \text{codim}(X, \mathbb{P}^N)$ and $\tilde{F}_i = \mathcal{O}_{\mathbb{P}^N}(-i)^{\oplus b_i}$.

(ii) *The cohomology $H^i(X, E(-p)) = 0$ for all $i > 0$ and $1 \leq p \leq \dim X$.*

(iii) *If $\pi : X \rightarrow \mathbb{P}^{\dim X}$ is a finite linear projection, then $\pi_*(E)$ is the trivial bundle.*

An ACM bundle E on a smooth hypersurface $X \subset \mathbb{P}^N$ comes equipped with a unique minimal resolution

$$0 \longrightarrow \tilde{F}_1 \xrightarrow{\Phi} \tilde{F}_0 \longrightarrow E \longrightarrow 0, \quad (1.1)$$

where \tilde{F}_1 and \tilde{F}_0 are sums of line bundles on \mathbb{P}^N . Here by *minimal*, we mean that no non-zero constant appears as an entry of the matrix Φ in (1.1). In this setting, an Ulrich bundle E is one which admits a *linear resolution*:

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^N}(-1)^M \xrightarrow{\Phi} \mathcal{O}_{\mathbb{P}^N}^M \longrightarrow E \longrightarrow 0.$$

It is known that $M = rd$, where r is the rank of E and d is the degree of the hypersurface. It follows from the resolution that E is 0-regular (in the sense of Castelnuovo and Mumford). It is also known that the dual E^\vee is $(d-1)$ -regular, and in particular $h^0(E^\vee) = 0$. We refer the reader to [5] and [11] for a quick introduction to Ulrich bundles, a discussion of their properties and their relevance.

Since any smooth complex hypersurface supports an Ulrich bundle [4], we may ask if these bundles are generic in the sense of the Franchetta-type conjectures. More precisely, one may ask:

Let $\mathcal{X} \rightarrow S$ be the universal family of smooth hypersurfaces of degree d in \mathbb{P}^N . Is there a vector bundle \mathcal{E} on \mathcal{X} which is flat over the base S such that its restriction to any smooth fiber X_s , $s \in S$, is an Ulrich bundle?

On the other hand, when a smooth family of varieties in a fixed ambient variety P , say $\mathcal{X} \subset S \times P$ is considered, it is known in various contexts that the behaviour of the generic fiber is related to that of P (see § 2, Proposition in [7] for a precise statement enunciating this principle). From this point of view, the Lefschetz theorems for Picard groups may be viewed as instances of the original Franchetta conjecture. Applying this principle to ACM bundles on a generic hypersurface (assuming that it holds) leads us to a version of Horrocks theorem.

For the sake of brevity, we adopt the following

Convention. Let $p : \mathcal{X} \rightarrow S$ be the universal family of smooth hypersurfaces of degree d in \mathbb{P}^N . Let $S' \rightarrow S$ be an étale map and \mathcal{E} on $\mathcal{X}' := \mathcal{X} \times_S S'$ be a vector bundle which is flat over S' such that for every $y \in S'$, the bundle \mathcal{E}_y on the hypersurface X_y is ACM. In what follows, the statement “ X is a general hypersurface of degree d in \mathbb{P}^N and E is an ACM bundle on X ” will mean that there exists an étale map $S' \rightarrow S$, a vector bundle \mathcal{E} as above, and a point $o \in S'$ such that $X = X_o$ and $E = \mathcal{E}_o$.

Indeed, results proved in the recent past support the above expectation. In the foundational article on this subject [6], it is shown that a general threefold of degree d in \mathbb{P}^4 supports a rank 2 Ulrich bundle if and only if $d \leq 5$. This was generalized to all rank 2 indecomposable ACM bundles (see [24, 29]). The corresponding statements for rank 3 bundles have been proved in [33, 30, 31]. These results establish the base case of a version of Franchetta's conjecture that has been proposed for low rank bundles in [30] where this is referred to as a *Noether-Lefschetz conjecture*. This conjecture in turn is inspired by a similar conjecture in [10] (see [20, 23, 32, 12] for progress related to this conjecture).

We prove the following generalization of these results.

Theorem 1.2. *There is no universal Ulrich bundle on smooth hypersurfaces of degree $d \geq 3$ and dimension $n \geq 4$.*

Theorem 1.2 follows from our main result below which uses a formalism introduced in [21] to prove the Noether-Lefschetz theorem for Picard groups.

Theorem 1.3. *Let X be a general hypersurface in \mathbb{P}^N of degree d and dimension at least 2, and let E be an ACM bundle on X . Let $n := \lfloor \frac{\dim X}{2} \rfloor$ (so that $\dim X = 2n$ or $2n + 1$). If the regularity of E and its dual E^\vee satisfy*

$$m(E), m(E^\vee) \leq nd - 2n - 2,$$

then any non-zero Hodge class $\zeta \in H^n(X, E \otimes \Omega_X^n)$ gives rise to a direct summand of E isomorphic to \mathcal{O}_X . In particular, the rank of the Hodge cohomology group $H^n(X, E \otimes \Omega_X^n)$ is bounded above by the rank of E .

As a consequence, we obtain the following result of which Theorem 1.2 is a special case.

Corollary 1.4. *Let X be a general hypersurface in \mathbb{P}^N of degree d and dimension at least 4, and let E be an ACM bundle on X . If E is initialized, i.e., $h^0(E(-1)) = 0 \neq h^0(E)$, and its regularity $m \leq nd - 2n - 2$, where $n := \lfloor \frac{\dim X}{2} \rfloor$, then E is isomorphic to a direct sum of line bundles.*

Remark 1.5. For an initialized ACM bundle E , we have $m(E) \geq 0$ and so $nd - 2n - 2 \geq 0$, or equivalently $d \geq 2 + \frac{2}{n}$. These degree bounds for odd-dimensional hypersurfaces are the same as the bounds that appear in the Noether-Lefschetz theory (cf. the work of Green [14] and Voisin (unpublished) on a conjecture of Griffiths and Harris generalizing the Noether-Lefschetz theorem).

Remark 1.6. Indecomposable ACM bundles on hypersurfaces are intimately related to Noether-Lefschetz theory in other ways as well. For example, a conjecture of Griffiths and Harris (see [15]) states that any curve C in a general hypersurface $X \subset \mathbb{P}^4$ is an intersection of the form $X \cap S$ for some surface $S \subset \mathbb{P}^4$, thus generalizing the usual Noether-Lefschetz theorem for surfaces in \mathbb{P}^3 . Voisin [34] constructs examples of curves disproving this conjecture. In [25], its authors provide a general framework using ACM bundles to construct a large class of codimension 2 subvarieties that do not arise as intersections and situates Voisin's examples amongst these.

Acknowledgements: The authors thank A. Beauville for a very useful e-mail exchange during the preparation of the manuscript. The authors thank the referee for not only a very careful reading of the manuscript but also for their comments and suggestions which

have provided a greater clarity for the results presented here. The second author was partially supported by a grant from the Simons Foundation.

2. PRELIMINARIES

The base field k is algebraically closed and its characteristic is zero.

Let $X \subset \mathbb{P} := \mathbb{P}^N$ be an irreducible smooth projective variety. Denote $W_1 := H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(1))$, and consider the dual of the Euler sequence

$$0 \longrightarrow \Omega_{\mathbb{P}}^1 \longrightarrow W_1^* \otimes \mathcal{O}_{\mathbb{P}}(-1) \longrightarrow \mathcal{O}_{\mathbb{P}} \longrightarrow 0.$$

It produces the Koszul complex

$$\begin{aligned} 0 \longrightarrow \Omega_{\mathbb{P}}^p \longrightarrow \wedge^p W_1^* \otimes \mathcal{O}_{\mathbb{P}}(-p) \longrightarrow \wedge^{p-1} W_1^* \otimes \mathcal{O}_{\mathbb{P}}(-p+1) \quad (2.1) \\ \dots \longrightarrow \dots \longrightarrow W_1^* \otimes \mathcal{O}_{\mathbb{P}}(-1) \longrightarrow \mathcal{O}_{\mathbb{P}} \longrightarrow 0 \end{aligned}$$

for every $p > 0$. Tensoring (2.1) with any vector bundle E on X and taking the corresponding long exact sequence of cohomologies, we obtain a coboundary map

$$\partial : H^0(X, E) \longrightarrow H^p(X, E \otimes \Omega_{\mathbb{P}}^p). \quad (2.2)$$

The following splitting criterion is a very mild generalization of an elegant result we learnt from the papers of Arrondo-Malaspina [2] and Arrondo-Tocino [3] and was the starting point for the present work.

Lemma 2.1. *Let $X \subset \mathbb{P}^N$ be a smooth projective variety with $\dim X = p + q$, and let E be a vector bundle on X with $H^0(X, E) \neq 0$. Assume that in the commutative diagram*

$$\begin{array}{ccc} H^0(E) \times H^0(E^\vee) & \xrightarrow{\varphi} & H^0(\mathcal{O}_X) \\ \downarrow \partial_1 \times \partial_2 & & \downarrow \partial \\ H^p(E \otimes \Omega_{\mathbb{P}}^p) \times H^q(E^\vee \otimes \Omega_{\mathbb{P}}^q) & \xrightarrow{\varphi_1} & H^{p+q}(\Omega_{\mathbb{P}|X}^{p+q}) \\ \downarrow \rho_1 \times \rho_2 & & \downarrow \rho \\ H^p(E \otimes \Omega_X^p) \times H^q(E^\vee \otimes \Omega_X^q) & \xrightarrow{\varphi_2} & H^{\dim X}(\omega_X) \end{array}$$

where ∂_1 , ∂_2 and ∂ are as in (2.2) while ρ_1 , ρ_2 and ρ are the restriction maps, the maps

- ∂_1 and ρ_1 are injective, and
- ∂_2 and ρ_2 are both surjective.

Then φ is a non-zero pairing. Consequently, the trivial line bundle is a direct summand of E .

Proof. Using contraction we have $(\Omega_X^p)^\vee \otimes \omega_X = \Omega_X^q$, and hence the map φ_2 in the diagram in the lemma is the Serre duality pairing. In particular, φ_2 is a perfect pairing.

Take a non-zero section $s \in H^0(E)$. Then $\rho_1 \circ \partial_1(s) \neq 0$, as both ρ_1 and ∂_1 are given to be injective. Since φ_2 is a perfect pairing, there exists a class $\xi \in H^q(E^\vee \otimes \Omega_X^q)$ such that $\varphi_2((\rho_1 \circ \partial_1(s)) \otimes \xi) \neq 0$. Finally, since the maps ∂_2 and ρ_2 are surjective, there is a section $s^\vee \in H^0(E^\vee)$ such that $\rho_2 \circ \partial_2(s^\vee) = \xi$. By the commutativity of the diagram we now have $\varphi(s \otimes s^\vee) \neq 0$.

Lemma 2.2 shows that the trivial line bundle is a direct summand of E . □

Lemma 2.2. *Let X be a smooth projective variety and E a vector bundle on X . Assume there is a non-zero pairing*

$$\theta : H^0(E) \times H^0(E^\vee) \longrightarrow H^0(\mathcal{O}_X).$$

Then $E \cong E' \oplus \mathcal{O}_X$.

Proof. Take $s \in H^0(E)$ and $s^\vee \in H^0(E^\vee)$ such that $\theta(s \otimes s^\vee) \neq 0$. These give nonzero homomorphisms $s^\vee : E \longrightarrow \mathcal{O}_X$ and $s : \mathcal{O}_X \longrightarrow E$ such that the composition of maps $s^\vee \circ s : \mathcal{O}_X \longrightarrow \mathcal{O}_X$ is non-zero, and hence $s^\vee \circ s$ is an isomorphism. This yields the required splitting. \square

Remark 2.3. In practice, finding reasonably general conditions under which *both* ρ_1 and ρ_2 satisfy the hypotheses in Lemma 2.1 can be challenging.

3. AN INFINITESIMAL LEFSCHETZ THEOREM FOR HODGE CLASSES OF AN ACM BUNDLE

As a prelude to our main result, we will establish, in Theorem 3.10, an *infinitesimal Lefschetz theorem* for Hodge classes of an ACM bundle on an even dimensional smooth projective hypersurface. The main tool here is a formalism, introduced by N. Mohan Kumar and V. Srinivas (unpublished), to prove the Noether-Lefschetz theorem for surfaces in \mathbb{P}^3 . An expository account of this can be found in [27]. Applications of this method are in [19], [25] and [28].

We start by setting up some notation that will be employed in this section.

As before, k is an algebraically closed field of characteristic zero. Set $\mathbb{P} := \mathbb{P}_k^N$ and $W_d = H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(d))$. Let $S := \mathbb{P}(W_d^*)$ denote the parameter space of all degree d hypersurfaces in \mathbb{P} . We have the short exact sequence

$$0 \longrightarrow \mathcal{V}_d \longrightarrow W_d \otimes \mathcal{O}_{\mathbb{P}} = H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(d)) \otimes \mathcal{O}_{\mathbb{P}} \longrightarrow \mathcal{O}_{\mathbb{P}}(d) \longrightarrow 0,$$

where \mathcal{V}_d is the kernel of the evaluation map $W_d \otimes \mathcal{O}_{\mathbb{P}} \longrightarrow \mathcal{O}_{\mathbb{P}}(d)$. Then

$$\mathcal{X} := \mathbb{P}(\mathcal{V}_d^*) \longrightarrow S \tag{3.1}$$

is the universal family of all degree d hypersurfaces. Let $X \subset \mathbb{P}$ be the smooth degree d hypersurface corresponding to a closed point $y \in S$.

We have the standard exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}} \longrightarrow \mathcal{O}_{\mathbb{P}}(d) \longrightarrow \mathcal{O}_X(d) \longrightarrow 0.$$

The corresponding long exact sequence of cohomologies gives

$$0 \longrightarrow k \longrightarrow W_d \longrightarrow V \longrightarrow 0,$$

where

$$V := H^0(X, \mathcal{O}_X(d)). \tag{3.2}$$

Then $V \cong T_{S,y}$, the Zariski tangent space to S at the point y . Consider the standard decomposition

$$A := \mathcal{O}_{S,y}/\mathfrak{m}_y^2 = k \oplus V^\vee,$$

so that $\Omega_A^1 \otimes k = H^0(\Omega_{\text{Spec } A}^1) \cong V^\vee$. We have the inclusion maps

$$\{y\} \hookrightarrow \text{Spec } A \hookrightarrow S;$$

note that $\text{Spec } A$ is the first order thickening of the smooth point $\{y\} \subset S$. Let

$$X_\epsilon := \text{Spec } A \times_S \mathcal{X}$$

denote the universal hypersurface over $\text{Spec } A$, where \mathcal{X} is defined in (3.1). So X_ϵ is the first order thickening of X as a subscheme of \mathcal{X} .

Set $\mathbb{P}_A := \mathbb{P} \times \text{Spec } A$; let $p : \mathbb{P}_A \rightarrow \mathbb{P}$ and $q : \mathbb{P}_A \rightarrow \text{Spec } A$ denote the two natural projections. Consider the conormal sheaf sequence for the inclusion map $\iota_\epsilon : X_\epsilon \hookrightarrow \mathbb{P}_A$

$$\mathcal{O}_{X_\epsilon}(-d) \rightarrow \Omega_{\mathbb{P}_A}^1 \otimes \mathcal{O}_{X_\epsilon} = p^* \Omega_{\mathbb{P}}^1 \otimes \mathcal{O}_{X_\epsilon} \oplus q^* \Omega_A^1 \otimes \mathcal{O}_{X_\epsilon} \rightarrow \Omega_{X_\epsilon}^1 \rightarrow 0.$$

Restricting this sequence to X yields the following:

$$\mathcal{O}_X(-d) \xrightarrow{(\alpha, \beta)} \Omega_{\mathbb{P}}^1 \otimes \mathcal{O}_X \oplus V^\vee \otimes \mathcal{O}_X \xrightarrow{(\gamma, \delta)} \Omega_{X_\epsilon}^1 \otimes \mathcal{O}_X \rightarrow 0.$$

Rewriting, we obtain the following

Lemma 3.1 (Mohan Kumar-Srinivas, [21]). *There is a commutative diagram*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X(-d) & \xrightarrow{\alpha} & \Omega_{\mathbb{P}}^1 \otimes \mathcal{O}_X & \longrightarrow & \Omega_X^1 \longrightarrow 0 \\ & & \downarrow -\beta & & \downarrow \gamma & & \parallel \\ 0 & \longrightarrow & V^\vee \otimes \mathcal{O}_X & \xrightarrow{\delta} & \Omega_{X_\epsilon}^1 \otimes \mathcal{O}_X & \longrightarrow & \Omega_X^1 \longrightarrow 0 \end{array} \quad (3.3)$$

where the top and bottom rows come from the inclusions $X \subset \mathbb{P}$ and $X \subset X_\epsilon$ respectively, while the homomorphism β is the dual of the evaluation map.

Taking the ℓ -th exterior power of (3.3) we obtain a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Omega_X^{\ell-1}(-d) & \longrightarrow & \Omega_{\mathbb{P}}^\ell \otimes \mathcal{O}_X & \longrightarrow & \Omega_X^\ell \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \Omega(\ell) & \longrightarrow & \Omega_{X_\epsilon}^\ell \otimes \mathcal{O}_X & \longrightarrow & \Omega_X^\ell \longrightarrow 0 \end{array} \quad (3.4)$$

(see [17, page 126, 5.16(d)]), where $\Omega(\ell)$ comes equipped with a decreasing filtration $F^* \Omega(\ell)$ for which

- $F^1(\Omega(\ell)) = \Omega(\ell)$,
- $F^{\ell+1}(\Omega(\ell)) = 0$, and
- $\text{gr}_{\mathbb{F}}^j \Omega(\ell) := F^j(\Omega(\ell)) / F^{j+1}(\Omega(\ell)) = \Lambda^j V^\vee \otimes \Omega_X^{\ell-j}$ for all $j \geq 1$.

Lemma 3.2. *For a smooth hypersurface $X \subset \mathbb{P}$ and an ACM bundle E on X ,*

$$H^n(X, E \otimes \Omega_X^{n-j}) = 0 = H^n(X, E(-d) \otimes \Omega_X^{n-j})$$

for all $j \geq 1$, where $n := \lfloor \dim X / 2 \rfloor$ (so $\dim X = 2n$ or $2n + 1$).

Proof. Associated to the cotangent bundle sequence

$$0 \longrightarrow \mathcal{O}_X(-d) \longrightarrow \Omega_{\mathbb{P}|X}^1 \longrightarrow \Omega_X^1 \longrightarrow 0 \quad (3.5)$$

we have the Koszul complex

$$\begin{aligned} 0 &\longrightarrow \mathcal{O}_X(-(n-j)d) \longrightarrow \Omega_{\mathbb{P}|X}^1(-(n-j-1)d) \longrightarrow \cdots \\ \cdots &\longrightarrow \Omega_{\mathbb{P}|X}^{n-j-1}(-d) \longrightarrow \Omega_{\mathbb{P}|X}^{n-j} \longrightarrow \Omega_X^{n-j} \longrightarrow 0. \end{aligned} \quad (3.6)$$

Tensoring (3.6) with E and taking the long exact sequence of cohomologies, we see that it is enough to prove the following for all $j \geq 1$:

- (a) $H^{2n-j}(X, E(-(n-j)d)) = 0$, and
- (b) $H^{n+a}(X, E \otimes \Omega_{\mathbb{P}|X}^{n-j-a}(-ad)) = 0$ for $0 \leq a < n-j$.

Statement (a) is clear from the fact that E is ACM, and $2n-j < \dim X = 2n$ or $2n+1$.

Statement (b) can be verified as follows: Tensoring the minimal resolution (1.1) of E with $\Omega_{\mathbb{P}}^{n-j-a}(-ad)$, we get that

$$0 \longrightarrow \tilde{F}_1 \otimes \Omega_{\mathbb{P}}^{n-j-a}(-ad) \xrightarrow{\Phi} \tilde{F}_0 \otimes \Omega_{\mathbb{P}}^{n-j-a}(-ad) \longrightarrow E \otimes \Omega_{\mathbb{P}}^{n-j-a}(-ad) \longrightarrow 0. \quad (3.7)$$

The long exact sequence of cohomologies associated to (3.7) produces the exact sequence

$$\begin{aligned} H^{n+a}(\mathbb{P}, \tilde{F}_0 \otimes \Omega_{\mathbb{P}}^{n-j-a}(-ad)) &\longrightarrow H^{n+a}(X, E \otimes \Omega_{\mathbb{P}|X}^{n-j-a}(-ad)) \\ &\longrightarrow H^{n+a+1}(\mathbb{P}, \tilde{F}_1 \otimes \Omega_{\mathbb{P}}^{n-j-a}(-ad)). \end{aligned} \quad (3.8)$$

Since $1 \leq n-j-a \leq n-1$ and $n+a < 2n-j < \dim X$, the two extreme terms in (3.8) vanish by Bott's formula. The proof of the second vanishing also follows along the same lines. \square

The following is a consequence of Lemma 3.2.

Corollary 3.3. *For a smooth, degree d hypersurface $X \subset \mathbb{P}^{2n+1}$ or $X \subset \mathbb{P}^{2n+2}$ and an ACM bundle E on X ,*

$$H^n(X, E \otimes \Omega(n)) = 0 = H^n(X, E(-d) \otimes \Omega(n)),$$

where $\Omega(n)$ is the filtered bundle in (3.4).

Proof. Consider the exact sequences

$$\begin{aligned} 0 \longrightarrow E \otimes F^{j+1}(\Omega(n)) &\longrightarrow E \otimes F^j(\Omega(n)) \longrightarrow E \otimes \text{gr}_{\mathbb{F}}^j \Omega \\ &= E \otimes \Lambda^j V^* \otimes \Omega_X^{n-j} \longrightarrow 0. \end{aligned} \quad (3.9)$$

In view of Lemma 3.2, the long exact sequence of cohomologies associated to (3.9) produces a surjection

$$H^n(X, E \otimes F^{j+1}(\Omega(n))) \rightarrow H^n(X, E \otimes F^j(\Omega(n))) \quad \forall j \geq 1.$$

Since $H^n(X, E \otimes F^n \Omega(n)) = H^n(X, E \otimes \Lambda^n V^* \otimes \mathcal{O}_X) = 0$, we obtain the first vanishing. The proof of the second vanishing is along the same lines. \square

Tensoring the Koszul complex

$$\begin{aligned} 0 \longrightarrow \mathcal{O}_X(-(n-1)d) &\longrightarrow \Omega_{\mathbb{P}|X}^1(-(n-2)d) \longrightarrow \cdots \\ \cdots &\longrightarrow \Omega_{\mathbb{P}|X}^{n-2}(-d) \longrightarrow \Omega_{\mathbb{P}|X}^{n-1} \longrightarrow \Omega_X^{n-1} \longrightarrow 0. \end{aligned} \quad (3.10)$$

with $E(-d)$ and E respectively, and taking the long exact sequence of cohomologies we get coboundary maps

$$H^{n+1}(X, E \otimes \Omega_X^{n-1}(-d)) \longrightarrow H^{2n}(X, E(-nd)), \quad (3.11)$$

$$H^{n+1}(X, E \otimes \Omega_X^{n-1}) \longrightarrow H^{2n}(X, E((-n+1)d)). \quad (3.12)$$

Lemma 3.4. *Let $X \subset \mathbb{P}^{2n+1}$, or $X \subset \mathbb{P}^{2n+2}$, be a smooth hypersurface of dimension at least two, and E an ACM bundle on X . Then the coboundary maps in (3.11) and (3.12) are injections.*

Proof. We prove the case when X has even dimension. The proof for odd dimensional hypersurfaces of dimension at least three is identical.

When $n = 1$, there is nothing to prove. So we assume that $n \geq 2$. We argue as in the proof of Lemma 3.2. From the Koszul complex (3.10) tensored with E and $E(-d)$ respectively, it is enough to show that

- (i) $H^{n+1+a}(X, E \otimes \Omega_{\mathbb{P}_1|X}^{n-1-a}(-ad)) = 0$ for $0 \leq a \leq n-2$, and
- (ii) $H^{n+1+a}(X, E \otimes \Omega_{\mathbb{P}_1|X}^{n-1-a}(-(a+1)d)) = 0$ for $0 \leq a \leq n-2$.

Tensoring the minimal resolution (1.1) of E with $\Omega_{\mathbb{P}}^{n-1-a}(-ad)$, and taking the corresponding long exact sequence of cohomologies, we get the exact sequence

$$\begin{aligned} H^{n+1+a}(\mathbb{P}, \tilde{F}_0 \otimes \Omega_{\mathbb{P}}^{n-1-a}(-ad)) &\longrightarrow H^{n+1+a}(X, E \otimes \Omega_{\mathbb{P}_1|X}^{n-1-a}(-ad)) \\ &\longrightarrow H^{n+2+a}(\mathbb{P}, \tilde{F}_1 \otimes \Omega_{\mathbb{P}}^{n-1-a}(-ad)). \end{aligned}$$

For the given values of a and n , the two extreme terms vanish by Bott's formula. This proves (i). The proof of (ii) is identical. \square

Proposition 3.5. *Let X be a smooth hypersurface and E an ACM vector bundle on X . Then the restriction map*

$$\rho : H^n(X, E \otimes \Omega_{\mathbb{P}}^n) \longrightarrow H^n(X, E \otimes \Omega_X^n)$$

is an injection when $\dim X = 2n$, and an isomorphism when $\dim X = 2n + 1$.

Proof. In view of the long exact sequence of cohomologies associated to the exact sequence

$$0 \longrightarrow \Omega_X^{n-1}(-d) \longrightarrow \Omega_{\mathbb{P}|X}^n \longrightarrow \Omega_X^n \longrightarrow 0,$$

to prove the injectivity of ρ , it suffices to show that $H^n(X, E \otimes \Omega_X^{n-1}(-d)) = 0$. Now this follows from Lemma 3.2.

The proof of surjectivity in the odd dimensional case is identical to the proof of Lemma 3.4. The vanishings in this case yield the injection

$$H^{n+1}(X, E \otimes \Omega_X^{n-1}(-d)) \hookrightarrow H^{2n}(X, E(-nd)).$$

However, since $\dim X = 2n + 1$, the cohomology on the right vanishes since E is ACM. This gives us the desired result. \square

In what follows, we shall say more about the image of the restriction map ρ in the above proposition in the even-dimensional case. For the sake of convenience, we make the following definition:

Definition 3.6. Let X be a smooth projective hypersurface of degree d , and $n = \lfloor \frac{\dim X}{2} \rfloor$. We say a vector bundle F on X satisfies Hypothesis (\star) if the natural homomorphism

$$H^0(X, F(nd - 2n - 2)) \otimes H^0(X, \mathcal{O}_X(d)) \longrightarrow H^0(X, F((n+1)d - 2n - 2)) \quad (\star)$$

is surjective.

Remark 3.7. Note that if F is $(nd - 2n - 2)$ -regular, then it satisfies Hypothesis (\star) .

Consider the composition of maps of sheaves

$$\Omega_X^{n-1} \otimes \mathcal{O}_X(-d) = \Omega_X^{n-1}(-d) \longrightarrow \Omega(n) \longrightarrow \Omega_X^{n-1} \otimes V^\vee = \mathrm{gr}_F^1 \Omega(n),$$

where the first map is the leftmost map in diagram (3.4) (set $\ell = n$ in (3.4)). This composition of maps coincides with the map $1 \otimes -\beta$ (see (3.3)). Let

$$\phi : H^{n+1}(X, E \otimes \Omega_X^{n-1}(-d)) \longrightarrow V^\vee \otimes H^{n+1}(X, E \otimes \Omega_X^{n-1}). \quad (3.13)$$

be the map of cohomologies induced by the above map of sheaves.

Lemma 3.8. *Let $X \subset \mathbb{P}^{2n+1}$ be a smooth hypersurface and E an ACM bundle on X such that the dual bundle E^\vee satisfies Hypothesis (\star) . Then the map ϕ in (3.13) is an injection.*

Proof. Using Lemma 3.4, we can identify ϕ with the restriction of the map

$$H^{2n}(X, E \otimes \mathcal{O}_X(-nd)) \longrightarrow V^\vee \otimes H^{2n}(X, E \otimes \mathcal{O}_X((-n+1)d)).$$

This is the dual of the cup product map

$$H^0(X, E^\vee(nd - 2n - 2)) \otimes H^0(X, \mathcal{O}_X(d)) \longrightarrow H^0(X, E^\vee((n+1)d - 2n - 2)).$$

By our hypothesis this cup product map is surjective. Hence the map ϕ is injective. \square

Proposition 3.9. *Let X and E be as in Lemma 3.8. Then there is an isomorphism*

$$H^n(X, E \otimes \Omega_{\mathbb{P}}^n) \cong H^n(X, E \otimes \Omega_{X_\epsilon}^n).$$

Proof. Consider the cohomology diagram associated to (3.4) for $\ell = n$:

$$\begin{array}{ccccc} H^n(X, E \otimes \Omega_X^{n-1}(-d)) & \longrightarrow & H^n(X, E \otimes \Omega_{\mathbb{P}}^n) & \longrightarrow & \\ & & \downarrow & & \\ H^n(X, E \otimes \Omega(n)) & \longrightarrow & H^n(X, E \otimes \Omega_{X_\epsilon}^n) & \longrightarrow & \\ \longrightarrow & H^n(X, E \otimes \Omega_X^n) & \longrightarrow & H^{n+1}(X, E \otimes \Omega_X^{n-1}(-d)) & \\ & \parallel & & \downarrow & \\ \longrightarrow & H^n(X, E \otimes \Omega_X^n) & \longrightarrow & H^{n+1}(X, E \otimes \Omega(n)). & \end{array} \quad (3.14)$$

From Lemma 3.2 and Corollary 3.3 it follows that

$$H^n(X, E \otimes \Omega_X^{n-1}(-d)) = H^n(X, E \otimes \Omega(n)) = 0.$$

Thus we see that the homomorphism

$$H^n(X, E \otimes \Omega_{\mathbb{P}}^n) \longrightarrow H^n(X, E \otimes \Omega_{X_\epsilon}^n)$$

is injective.

To prove surjectivity, it suffices to show that the right most vertical homomorphism in (3.14), i.e., the map $H^{n+1}(X, E \otimes \Omega_X^{n-1}(-d)) \longrightarrow H^{n+1}(X, E \otimes \Omega(n))$ is injective. By Lemma 3.8, the map ϕ , which is the composition of maps

$$H^{n+1}(X, E \otimes \Omega_X^{n-1}(-d)) \longrightarrow H^{n+1}(X, E \otimes \Omega(n)) \longrightarrow V^\vee \otimes H^{n+1}(X, E \otimes \Omega_X^{n-1}),$$

is injective. Therefore, the *first* map $H^{n+1}(X, E \otimes \Omega_X^{n-1}(-d)) \longrightarrow H^{n+1}(X, E \otimes \Omega(n))$ is injective. \square

Theorem 3.10 (The infinitesimal Lefschetz theorem for Hodge classes). *Let $X \subset \mathbb{P}^{2n+1}$ be a smooth hypersurface and E an ACM bundle on X such that its dual satisfies the Hypothesis (\star) . Then there exists an exact sequence*

$$0 \longrightarrow H^n(X, E \otimes \Omega_{X_\epsilon}^n|_X) \longrightarrow H^n(X, E \otimes \Omega_X^n) \xrightarrow{\kappa} V^\vee \otimes H^{n+1}(X, E \otimes \Omega_X^{n-1}).$$

Proof. The map κ is just the composition of maps

$$H^n(X, E \otimes \Omega_X^n) \longrightarrow H^{n+1}(X, E \otimes \Omega(n)) \longrightarrow V^\vee \otimes H^{n+1}(X, E \otimes \Omega_X^{n-1}).$$

The theorem now follows from the identification in Proposition 3.9. \square

Remark 3.11. The homomorphism κ in Theorem 3.10 is in fact the Kodaira-Spencer map

$$\begin{array}{ccc} H^n(X, E \otimes \Omega_X^n) & \longrightarrow & \text{Hom}(T_{S,o}, H^{n+1}(X, E \otimes \Omega_X^{n-1})) \\ \xi & \longmapsto & (\frac{\partial}{\partial x} \longmapsto \frac{\partial}{\partial x}(\xi)). \end{array} \quad (3.15)$$

Hence, if X is general fiber in the universal family of degree d hypersurfaces $\mathcal{X} \rightarrow S$, so that ξ is the restriction of a section in $H^0(S', R^n p_*(\mathcal{E} \otimes \Omega_{\mathcal{X}'/S'}^n))$, then $\xi \mapsto 0$ under the Kodaira-Spencer map. In particular, when $E \cong \mathcal{O}_X$, the cohomology group $H^n(X, \Omega_{X_\epsilon}^n|_X)$ can be identified with the subspace of cohomology classes of type (n, n) on X which deform infinitesimally, i.e., they continue to be of type (n, n) under infinitesimal deformations. This statement is the usual *infinitesimal Noether-Lefschetz theorem* (see [27]).

4. PROOF OF THEOREM 1.3

We now apply the splitting criterion in Lemma 2.1 to the case of interest to us where E is an ACM bundle on a general hypersurface of dimension at least *two*.

Lemma 4.1. *Let $X \subset \mathbb{P}^N$ be an irreducible smooth projective variety and let F be a vector bundle on X . Set $n := \lfloor \frac{\dim X}{2} \rfloor$, so that $\dim X$ is either $2n$ or $2n + 1$. Then the map*

$$\partial : H^0(X, F) \longrightarrow H^n(X, F \otimes \Omega_{\mathbb{P}}^n)$$

in (2.2) is surjective, if

$$H^i(X, F(-i)) = 0$$

for all $1 \leq i \leq n$.

Proof. Tensor the Koszul complex (2.1) with F . Breaking up the resulting exact sequence into short exact sequences, we get

$$0 \longrightarrow F \otimes \Omega_{\mathbb{P}}^{k+1} \longrightarrow F \otimes \wedge^{k+1} W_1^\vee \otimes \mathcal{O}_{\mathbb{P}}(-k-1) \longrightarrow F \otimes \Omega_{\mathbb{P}}^k \longrightarrow 0$$

for all $0 \leq k \leq n-1$. Consider the corresponding long exact sequence of cohomologies. Incorporating the hypotheses, a surjection

$$\tilde{\partial}_k : H^k(X, F \otimes \Omega_{\mathbb{P}}^k) \longrightarrow H^{k+1}(X, F \otimes \Omega_{\mathbb{P}}^{k+1})$$

is obtained for every $0 \leq k \leq n-1$. Therefore, the composition of homomorphisms

$$\partial = \tilde{\partial}_{n-1} \circ \cdots \circ \tilde{\partial}_0 : H^0(X, F) \longrightarrow H^n(X, F \otimes \Omega_{\mathbb{P}}^n)$$

is surjective. \square

Corollary 4.2. *In the statement of Lemma 4.1, if we assume in addition that*

$$H^i(X, F(-i-1)) = 0, \quad 0 \leq i < n,$$

then the map ∂ is an isomorphism.

4.1. The case of even-dimensional hypersurfaces.

Theorem 4.3. *Let $X \subset \mathbb{P}^{2n+1}$ be a general hypersurface of degree d and dimension at least 2, and let E be an ACM bundle on X such that both E and its dual E^\vee satisfy Hypothesis (\star) . Then any non-zero class $\zeta \in H^n(X, E \otimes \Omega_X^n)$ yields a direct summand of E isomorphic to \mathcal{O}_X .*

Proof. We recall our convention in §1. Let $p : \mathcal{X} \rightarrow S$ be the universal family of smooth hypersurfaces of degree d in $\mathbb{P} = \mathbb{P}^{2n+1}$, and let $q : \mathcal{X} \rightarrow \mathbb{P}$ be the natural projection. Then there exists an étale map $S' \rightarrow S$ and an ACM bundle \mathcal{E} on $\mathcal{X}' := \mathcal{X} \times_S S'$ which is flat over S' such that for every $y \in S'$, the ACM bundle \mathcal{E}_y on the hypersurface X_y satisfies the hypothesis in Lemma 3.8.

Let $o \in S'$ be a point such that $X = X_0$. Consider the diagram

$$\begin{array}{ccccc} X & \hookrightarrow & X_\epsilon & \hookrightarrow & \mathcal{X}' \\ \downarrow & & \downarrow & & \downarrow \\ \text{Spec } \mathbb{C} & \hookrightarrow & \text{Spec } A & \hookrightarrow & S' \end{array} \quad (4.1)$$

With an abuse of notation, all the vertical maps in (4.1) are denoted by p . By shrinking S' if necessary, we have

$$H^0(\mathcal{X}', \mathcal{E}) \cong H^0(S', p_*\mathcal{E}) \cong (p_*\mathcal{E})_o \cong H^0(X, E) \xrightarrow{\partial_1} H^n(X, E \otimes \Omega_{\mathbb{P}}^n) \xrightarrow{\rho_1} H^n(X, E \otimes \Omega_X^n).$$

Here ∂_1 is a surjection by Lemma 4.1 and ρ_1 is an injection by Proposition 3.5. A quick note on the isomorphisms above: the first isomorphism follows from a Leray spectral sequence argument, whereas the second one follows since $p_*\mathcal{E}$ is locally trivial and we are allowed to shrink S' if necessary. The third isomorphism follows from the proper base change formula.

Since E is a bundle on a general hypersurface X , once again, by the proper base change formula, we have the identification

$$(R^n p_*(\mathcal{E}^\vee \otimes \Omega_{\mathcal{X}'/S'}^n))_o \cong H^n(X, E \otimes \Omega_X^n),$$

under which the Hodge class ζ comes from a global class. Consequently (see Remark 3.11), it is in the kernel of

$$\kappa : H^n(X, E \otimes \Omega_X^n) \rightarrow V^\vee \otimes H^{n+1}(X, E \otimes \Omega_X^{n-1}).$$

Since E^\vee satisfies Hypothesis (\star) , by Theorem 3.10 and Proposition 3.9,

$$\zeta \in H^n(X, E \otimes \Omega_{\mathbb{P}}^n).$$

Let

$$\tilde{s} \in H^0(\mathcal{X}', \mathcal{E}), \quad (4.2)$$

with

$$s \in H^0(X, E) \quad (4.3)$$

its restriction to the fiber X , such that $\partial_1(s) = \zeta$. Then, via the perfect pairing φ_2 in Lemma 2.1,

$$H^n(X, E \otimes \Omega_X^n) \times H^n(X, E^\vee \otimes \Omega_X^n) \xrightarrow{\varphi_2} H^{2n}(X, \omega_X)$$

its image $\rho_1(\partial_1 s) \in H^n(X, E \otimes \Omega_X^n)$ defines a non-zero map

$$\zeta : H^n(X, E^\vee \otimes \Omega_X^n) \longrightarrow \mathbb{C}, \quad \xi \longmapsto \varphi_2(\zeta, \xi) \in \mathbb{C}.$$

This perfect pairing is obtained by taking the fiber at the point $o \in S'$ of the perfect pairing (= Serre duality in families)

$$R^n p_*(\mathcal{E} \otimes \Omega_{X'/S'}^n) \times R^n p_*(\mathcal{E}^\vee \otimes \Omega_{X'/S'}^n) \longrightarrow R^{2n} p_* \omega_{X'/S'} \xrightarrow{tr} \mathcal{O}_{S'}.$$

Here tr is trace map.

Thus, we have a homomorphism

$$\tilde{\zeta} : R^n p_*(\mathcal{E}^\vee \otimes \Omega_{X'/S'}^n) \longrightarrow \tilde{\mathbb{C}}.$$

This $\tilde{\zeta}$ is non-zero at the point $o \in S'$. Consequently, there is a neighbourhood $U \subset S'$ of o on which $\tilde{\zeta}$ is non-zero everywhere. This implies that there is a non-zero section

$$\tilde{\xi} \in H^0(U, R^n p_*(\mathcal{E}^\vee \otimes \Omega_{X'/S'}^n))$$

such that $(\rho_1(\partial_1 \tilde{s}), \tilde{\xi})$ is a non-zero section of $\tilde{\mathbb{C}}$ (see (4.2) for \tilde{s}). Since $\tilde{\xi}$ is defined over a neighbourhood U , this implies that it *deforms infinitesimally*, and hence at the point $o \in U$ we have

$$\xi_o := \tilde{\xi}(0) \in \text{kernel}(\kappa).$$

As E satisfies Hypothesis (\star) , by Lemma 4.1, and the identifications in Theorem 3.10 and Proposition 3.9, we have

$$H^0(X, E^\vee) \xrightarrow{\partial_2} H^n(X, E^\vee \otimes \Omega_{\mathbb{P}^n}^n) \xrightarrow{\rho_2} H^n(X, E^\vee \otimes \Omega_X^n).$$

Thus, there exists $t \in H^0(X, E^\vee)$ such that $\xi_o = \rho_2(\partial_2(t))$, and in particular we see that $(s, t) \neq 0$ (see (4.3) for s) under the pairing

$$H^0(X, E) \times H^0(X, E^\vee) \longrightarrow H^0(X, \mathcal{O}_X).$$

This implies that $E \cong E' \oplus \mathcal{O}_X$. □

4.2. Odd dimensional hypersurface. We begin with the following lemma.

Lemma 4.4. *Let X be a smooth hypersurface in \mathbb{P}^{2n+2} . Suppose that E is an ACM bundle with a section s such that its image*

$$\xi := \rho \circ \partial(s) \in H^n(X, E \otimes \Omega_X^n)$$

is non-zero (see Lemma 4.1 and Proposition 3.5 for ∂ and ρ respectively). Then there is a smooth hyperplane section $Y \subset X$ with defining polynomial $\ell \in H^0(X, \mathcal{O}_X(1))$ such that the image of the restriction s_Y in $H^n(Y, E \otimes \Omega_Y^n)$, denoted by ξ_Y , is also non-zero.

Proof. First note that we have a series of isomorphisms

$$H^i(X, E \otimes \Omega_{\mathbb{P}^n}^i) \cong H^{i+1}(X, E \otimes \Omega_{\mathbb{P}^n}^{i+1}), \quad 1 \leq i \leq n-1.$$

Let Y be a general hyperplane section and let $E_1 := E \otimes \mathcal{O}_Y$. We have a commutative diagram obtained from the Euler sequence

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
\rightarrow & W_1 \otimes H^0(X, E(-2)) & \rightarrow & H^0(X, E(-1)) & \rightarrow & H^1(X, E(-1) \otimes \Omega_{\mathbb{P}^{2n+2}}^1) & \rightarrow 0 \\
& & \downarrow & & \downarrow \times \ell & & \downarrow \\
\rightarrow & W_1 \otimes H^0(X, E(-1)) & \rightarrow & H^0(X, E) & \rightarrow & H^1(X, E \otimes \Omega_{\mathbb{P}^{2n+2}}^1) & \rightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
\rightarrow & W_1 \otimes H^0(X, E_1(-1)) & \rightarrow & H^0(Y, E_1) & \rightarrow & H^1(Y, E \otimes \Omega_{\mathbb{P}^{2n+2}}^1 \otimes \mathcal{O}_Y) & \rightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0
\end{array}$$

Further assume that Y is such that the restriction $s_Y \neq 0$. Suppose that

$$s_Y \mapsto \xi_Y = 0$$

in the bottom row. This means that s_Y is the restriction of

$$\sum_i x_i s_i \in W_1 \otimes H^0(X, E(-1)),$$

where x_i are the homogeneous coordinates. Thus, we see that

$$s - \sum_i x_i s_i = \ell \cdot t$$

for some $t \in H^0(X, E(-1))$, where s is the section in the statement of the lemma. In particular, $s = \sum_i x_i s'_i$, and hence the image of s in $H^1(X, E \otimes \Omega_{\mathbb{P}^{2n+2}}^1)$ is zero. This is a contradiction. \square

Theorem 4.5. *Let $X \subset \mathbb{P}^{2n+2}$ be a general hypersurface of degree d and dimension at least 3, and let E be an ACM bundle on X such that both E and its dual E^\vee satisfies Hypothesis (\star) . Then any non-zero class $\xi \in H^n(X, E \otimes \Omega_X^n)$ yields a direct summand of E isomorphic to \mathcal{O}_X .*

Proof. Since $X \subset \mathbb{P}^{2n+2}$ is a general hypersurface of degree d , and dimension at least 3, a general hyperplane section of X will yield a general hypersurface Y of degree d in \mathbb{P}^{2n+1} . As in the statement, E and E^\vee are ACM vector bundles on X satisfying Hypothesis (\star) . Then the restrictions $E_1 := E|_Y$ and E_1^\vee are also ACM bundles on Y satisfying Hypothesis (\star) .

Consider the sequence of maps

$$H^0(X, E) \xrightarrow{\partial} H^n(X, E \otimes \Omega_{\mathbb{P}^{2n+2}}^n) \xrightarrow{\rho} H^n(X, E \otimes \Omega_X^n).$$

By Proposition 3.5, the map ρ is an isomorphism, and by Lemma 4.1, the map ∂ is a surjection. Hence there exists a non-zero section $s \in H^0(X, E)$ such that

$$\rho \circ \partial(s) = \xi.$$

Furthermore, by Lemma 4.4, ξ_Y is a non-zero Hodge class and consequently, by Theorem 4.3, the vector bundle E_1 splits off a trivial rank one summand.

We claim that this splitting lifts to X as well. To prove this, note that any map $\mathcal{O}_Y \rightarrow E_1$ (respectively, $E_1 \rightarrow \mathcal{O}_Y$) lifts to a map $\mathcal{O}_X \rightarrow E$ (respectively, $E \rightarrow \mathcal{O}_X$) because E and E^\vee are ACM (the restriction homomorphism $H^0(X, F) \rightarrow H^0(Y, F_Y)$ is surjective

for any ACM bundle F on X). Now the composition of maps $\mathcal{O}_X \rightarrow E \rightarrow \mathcal{O}_X$ is nowhere zero since its restriction to Y is nowhere zero. Hence E has a direct summand isomorphic to \mathcal{O}_X . \square

4.3. Proofs of Theorems.

Proof of Theorem 1.2. The proof is by contradiction. Let X be a general degree d hypersurface of dimension at least four and suppose that X supports an Ulrich bundle E . Since E is Ulrich, we have that

- (i) E is initialized and 0-regular,
- (ii) E^\vee is $(d-1)$ -regular, and
- (iii) $h^0(E^\vee) = 0$.

It follows from these properties, and the exact sequence

$$0 \rightarrow E(-1) \rightarrow E \rightarrow E|_Y \rightarrow 0,$$

that the restriction of an Ulrich bundle on X to a hyperplane section $Y \subset X$ is also Ulrich (see § 3, [5]). Hence, without loss of generality, we may assume that X is even-dimensional, namely $\dim X = 2n \geq 4$.

Since E is initialized, by Corollary 4.2, there exists a non-zero Hodge cycle $\xi \in H^n(X, E \otimes \Omega_X^n)$, and by Serre duality a non-zero cycle $\xi^\perp \in H^n(X, E^\vee \otimes \Omega_X^n)$. The arguments in the proof of Theorem 4.3 show that ξ^\perp deforms infinitesimally and hence $\xi^\perp \in \ker(\kappa)$.

Since E is 0-regular, this means that E satisfies Hypothesis (\star) and so by Theorem 3.10 (applied to the (ACM) bundle E^\vee) and the surjectivity of the map ∂ (Lemma 4.1), we see that $h^0(E^\vee) \neq 0$. This is a contradiction. \square

Proof of Theorem 1.3. By definition, if a bundle is a -regular, then it is also $(a+i)$ -regular for any $i \geq 0$. Therefore, E and E^\vee satisfy Hypothesis (\star) . The result now follows from Theorem 4.3 and Theorem 4.5. \square

Proof of Corollary 1.4. We note that the dual of any initialized ACM bundle is $(d-1)$ -regular, and so both E and E^\vee satisfy Hypothesis (\star) . Furthermore, we have an isomorphism

$$H^0(X, E) \cong H^n(X, E \otimes \Omega_{\mathbb{P}}^n),$$

where $n = \lfloor \dim X/2 \rfloor$ (Corollary 4.2). Hence E has non-zero (n, n) Hodge cycles. Arguing as above, we see that $E \cong E' \oplus \mathcal{O}_X^{h^0(E)}$.

Let $a \geq 0$ be such that $E'_a := E'(a)$ is initialized. Then $m(E'_a) = m(E') - a \leq m(E)$. Hence E'_a satisfies the hypothesis and thus we have $E'_a \cong E'' \oplus \mathcal{O}_X^{h^0(E'_a)}$, or equivalently, E' has a direct summand of the form $\mathcal{O}_X(-a)^{h^0(E'_a)}$.

Continuing in this fashion, we see that E decomposes into a sum of line bundles. \square

REFERENCES

- [1] E. Arbarello and M. Cornalba, The Picard groups of the moduli spaces of curves, *Topology* **26** (1987), 153–171.

- [2] E. Arrondo and F. Malaspina, Cohomological characterization of vector bundles on Grassmannians of lines, *J. Algebra* **323** (2010), 1098–1106.
- [3] E. Arrondo and A. Tocino, Cohomological characterization of Universal Bundles on $\mathbb{G}(1, n)$, *J. Algebra* **540** (2019), 206–233.
- [4] J. Herzog, B. Ulrich and J. Backelin, Linear maximal Cohen-Macaulay modules over strict complete intersections, *J. Pure Appl. Alg.* **71** (1991), 187–202.
- [5] A. Beauville, An introduction to Ulrich bundles, *Eur. J. Math.* **4**, no. 1 (2018), 26–36.
- [6] A. Beauville, Determinantal hypersurfaces, *Michigan Math. J.* **48** (2000), 39–64.
- [7] A. Beauville, A remark on the generalized Franchetta conjecture for K3 surfaces, *Math Zeit.* (to appear).
- [8] A. Beauville and C. Voisin, On the Chow ring of a K3 surface, *J. Algebraic Geom.* **13** (2004), 417–426.
- [9] N. Bergeron and Z. Li, Tautological classes on moduli spaces of hyper-Kähler manifolds, *Duke Math. J.* **168** (2019), 1179–1230.
- [10] R.-O. Buchweitz, G.-M. Greuel and F.-O. Schreyer, Cohen-Macaulay modules on hypersurface singularities II, *Invent. Math.* **88** (1987) 165–182.
- [11] Emre Coskun, A survey of Ulrich bundles, *Analytic and algebraic geometry*, 85–106, Hindustan Book Agency, New Delhi, 2017.
- [12] D. Erman, Matrix factorizations of generic polynomials, arXiv preprint.
- [13] A. Franchetta, Sulle serie lineari razionalmente determinate sulla curva a moduli generali di dato genere, *Matematiche (Catania)* **9** (1954), 126–147.
- [14] Mark Green, Griffiths' infinitesimal invariant and the Abel-Jacobi map, *J. Differential Geom.* **29** (1989), no. 3, 545–555.
- [15] P. Griffiths and J. Harris, On the Noether-Lefschetz theorem and some remarks on codimension-two cycles, *Math. Ann.* **271** (1985), 31–51.
- [16] J. L. Harer, The second homology group of the mapping class group of an orientable surface, *Invent. Math.* **72** (1983), 221–239.
- [17] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, No. 52. Springer-Verlag, 1977.
- [18] G. Horrocks, Vector bundles on the punctured spectrum of a local ring, *Proc. London Math. Soc.* **14** (1964), 689–713.
- [19] K. Joshi, A Noether-Lefschetz theorem and applications, *J. Algebraic Geom.* **4** (1995), 105–135.
- [20] H. Kleppe, Deformation of schemes defined by vanishing of Pfaffians, *J. Algebra* **53** (1978), 84–92.
- [21] N. Mohan Kumar and V. Srinivas, The Noether-Lefschetz theorem, Unpublished notes.
- [22] N. Mohan Kumar, A. P. Rao and G. V. Ravindra, Generators for vector bundles on generic hypersurfaces, *Math. Res. Lett.* **14** (2007), 649–655.
- [23] N. Mohan Kumar, A.P. Rao and G.V. Ravindra, Arithmetically Cohen-Macaulay bundles on hypersurfaces, *Comment. Math. Helv.* **82** (2007), 829–843.
- [24] N. Mohan Kumar, A.P. Rao, and G.V. Ravindra, Arithmetically Cohen-Macaulay bundles on three dimensional hypersurfaces, *Int. Math. Res. Not.* (2007) no. 8, Art. ID rnm025, 11 pp.
- [25] N. Mohan Kumar, A. P. Rao and G. V. Ravindra, On codimension two subvarieties in hypersurfaces, *Motives and algebraic cycles*, 167–174, Fields Inst. Commun., 56, Amer. Math. Soc., Providence, RI, 2009.
- [26] K. O'Grady, Moduli of sheaves and the Chow group of K3 surfaces, *J. Math. Pures Appl.* **100** (2013), 701–718.
- [27] G. V. Ravindra, The Noether-Lefschetz theorem via vanishing of coherent cohomology, *Canad. Math. Bull.* **51** (2008), 283–290.
- [28] G. V. Ravindra and V. Srinivas, The Noether-Lefschetz theorem for the divisor class group, *J. Algebra* **322** (2009), 3373–3391.
- [29] G.V. Ravindra, Curves on threefolds and a conjecture of Griffiths-Harris, *Math. Ann.* **345** (2009), 731–748.
- [30] G.V. Ravindra, and A. Tripathi, Rank 3 ACM bundles on general hypersurfaces in \mathbb{P}^5 , *Adv. Math.* **355** (2019), 106780.
- [31] G.V. Ravindra and A. Tripathi, On the base case of a conjecture on ACM bundles over hypersurfaces, *Geom. Dedicata* **216** (2022), issue 5, 10pp.

- [32] A. Tripathi, Splitting of low-rank ACM bundles on hypersurfaces of high dimension, *Comm. Alg.* **44** (2016), 1011–1017.
- [33] A. Tripathi, Rank 3 arithmetically Cohen-Macaulay bundles over hypersurfaces, *J. Algebra* **478** (2017), 1–11.
- [34] C. Voisin, Sur une conjecture de Griffiths et Harris, *Algebraic curves and projective geometry* (Trento, 1988), 270–275, Lecture Notes in Math., 1389, Springer, Berlin, 1989.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD,
MUMBAI 400005, INDIA

E-mail address: `indranil@math.tifr.res.in`

UNIVERSITY OF MISSOURI – ST. LOUIS, ST. LOUIS, MO 63121, USA

E-mail address: `girivarur@umsl.edu`