

Fourier Transform of Bernstein-Bézier Polynomials

Charles. K. Chui^{1*}, Tian-Xiao He², and Qingtang Jiang³

¹ Department of Mathematics and Computer Science

University of Missouri-St. Louis, St. Louis, MO 63121, USA

and

Department of Statistics, Stanford University, Stanford, CA 94305, USA

² Department of Mathematics and Computer Science

Illinois Wesleyan University, Bloomington, IL 61702-2900, USA

³ Department of Mathematics and Computer Science

University of Missouri-St. Louis, St. Louis, MO 63121, USA

Abstract

Explicit formulae, in terms of Bernstein-Bézier coefficients, of the Fourier transform of bivariate polynomials on a triangle and univariate polynomials on an interval are derived in this paper. Examples are given and discussed to illustrate the general theory. Finally, this consideration is related to the study of refinement masks of spline function vectors.

AMS Subject Classification: 42C40, 42C15, 41A15.

Key Words and Phrases: Fourier transform, Bernstein-Bézier polynomials, minimal support, quasi-minimal support, refinement

1 Introduction

The objective of this paper is to present a compact formula of the Fourier transform of bivariate polynomials on a triangle and univariate polynomials on a bounded interval in terms of their Bernstein-Bézier (BB) coefficients. Of course the BB coefficients are formulated, as usual, in terms of the Barycentric coordinates, as opposed to the Cartesian coordinates $\mathbf{x} = (x_1, x_2)$ for $\mathbf{x} \in \mathbb{R}^2$ or $x \in \mathbb{R}$. We will focus on the bivariate setting and only consider the univariate formulae as simple consequences. In this regard, although our method of derivation can be

*Research supported by ARO Grant #W911NF-04-1-0298.

extended to multivariate polynomials on simplexes, we have decided to present the detailed derivation only for bivariate polynomials, since the motivation of this research is the study of subdivision masks for parametric spline curves and surfaces.

Let $P_n(\mathbf{x})$, $\mathbf{x} \in \mathbb{R}^2$, $\mathbf{x} = (x_1, x_2)$, be a Bernstein-Bézier polynomial on a triangle $\triangle A_1 A_2 A_3$ with vertices $A_i = (a_i, b_i)$ ($i = 1, 2, 3$). We write $P_n(\mathbf{x})$ in terms of the Barycentric coordinates (u, v, w) of $\triangle A_1 A_2 A_3$ as follows.

$$P_n(\mathbf{x}) \equiv P_n(x_1, x_2) = \sum_{0 \leq j, k, l \leq n, j+k+l=n} a_{j,k,l} \frac{n!}{j!k!l!} u^j v^k w^l, \quad (1.1)$$

where (u, v, w) is the Barycentric coordinates of $\mathbf{x} = (x_1, x_2) \in \triangle A_1 A_2 A_3$; i.e., $(x_1, x_2) = (a_1 u + a_2 v + a_3 w, b_1 u + b_2 v + b_3 w)$, $0 \leq u, v, w \leq 1$, $u + v + w = 1$.

We need the following notation. For $(m, s) \in \{(1, 2), (1, 3), (2, 3)\}$, the forward-backward operator $\triangle_{m,s}$ and backward-forward operator $\nabla_{m,s}$ defined on sequences $\{a_{j,k,l}\}_{j,k,l}$ of multi-indices j, k, l are given by:

$$\begin{aligned} \triangle_{12} a_{j,k,l} &:= a_{j+1,k-1,l} - a_{j,k,l}, & \nabla_{12} a_{j,k,l} &:= a_{j,k,l} - a_{j-1,k+1,l}, \\ \triangle_{13} a_{j,k,l} &:= a_{j+1,k,l-1} - a_{j,k,l}, & \nabla_{13} a_{j,k,l} &:= a_{j,k,l} - a_{j-1,k,l+1}, \\ \triangle_{23} a_{j,k,l} &:= a_{j,k+1,l-1} - a_{j,k,l}, & \nabla_{23} a_{j,k,l} &:= a_{j,k,l} - a_{j,k-1,l+1}. \end{aligned} \quad (1.2)$$

In addition, we set $\triangle_{m,s}^k := \triangle_{m,s}(\triangle_{m,s}^{k-1})$ and $\nabla_{m,s}^k := \nabla_{m,s}(\nabla_{m,s}^{k-1})$ for $k = 1, 2, \dots$.

For a triangle $T = \triangle A_1 A_2 A_3$, we use V_T to denote its area, given by

$$V_T := \frac{1}{2} \left| \det \left(\begin{bmatrix} 1 & 1 & 1 \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{bmatrix} \right) \right|.$$

The main result of this paper can be stated as follows.

Theorem 1.1 *The Fourier transform of $P_n(\mathbf{x})$ as in (1.1) over a triangle $\triangle A_1 A_2 A_3$ has the explicit formulation:*

$$\begin{aligned} \widehat{P}_n(\xi) &:= \int_{\triangle A_1 A_2 A_3} P_n(\mathbf{x}) e^{-i\xi \cdot \mathbf{x}} d\mathbf{x} \\ &= 2V_T \sum_{\ell=0}^n \sum_{k=0}^{n-\ell} (-1)^{k+\ell} \frac{n!}{(n-k-\ell)!} \frac{1}{\gamma^{\ell+1}} \\ &\quad \times \left\{ \frac{1}{\alpha^{k+1}} \left(\nabla_{12}^k \nabla_{23}^\ell a_{n-\ell,0} e^{-i(a_1 \xi_1 + b_1 \xi_2)} - \triangle_{12}^k \nabla_{23}^\ell a_{0,n,0} e^{-i(a_2 \xi_1 + b_2 \xi_2)} \right) \right. \\ &\quad \left. - \frac{1}{\beta^{k+1}} \left(\nabla_{13}^k \triangle_{23}^\ell a_{n-\ell,0,\ell} e^{-i(a_1 \xi_1 + b_1 \xi_2)} - \triangle_{13}^k \triangle_{23}^\ell a_{0,0,n} e^{-i(a_3 \xi_1 + b_3 \xi_2)} \right) \right\}, \end{aligned} \quad (1.3)$$

where $\xi = (\xi_1, \xi_2)$, and $\alpha := i((a_2 - a_1)\xi_1 + (b_2 - b_1)\xi_2)$, $\beta := i((a_3 - a_1)\xi_1 + (b_3 - b_1)\xi_2)$, $\gamma := i((a_3 - a_2)\xi_1 + (b_3 - b_2)\xi_2)$.

The above formulation is valid for univariate polynomials by setting $w = 0$, namely:

$$p_n(x) = \sum_{0 \leq j, k \leq n, j+k=n} b_{j,k} \frac{n!}{j!k!} u^j v^k, \quad (1.4)$$

with $u = (x - b)/(a - b)$, $v = (x - a)/(b - a)$, and $x \in [a, b]$. By using Δ and ∇ to denote the forward-backward and backward-forward operators:

$$\Delta b_{j,k} := b_{j+1,k-1} - b_{j,k}, \quad \nabla b_{j,k} := b_{j,k} - b_{j-1,k+1}, \quad (1.5)$$

and $\Delta^k := \Delta(\Delta^{k-1})$ and $\nabla^k := \nabla(\nabla^{k-1})$, for $k = 1, 2, \dots$. Then the formulation of the Fourier transform of univariate polynomials is an immediate consequence of Theorem 1.1, as follows.

Corollary 1.2 *The Fourier transform of $p_n(x)$ in (1.4) over a bounded interval $[a, b]$ has the explicit formulation:*

$$\begin{aligned} \hat{p}_n(\xi) &:= \int_a^b p_n(x) e^{-ix\xi} dx \\ &= (b - a) e^{-ib\xi} \sum_{k=0}^n (-1)^k \frac{n!}{(n-k)!} \frac{1}{(i(b-a)\xi)^{k+1}} \left(\nabla^k b_{n,0} - \Delta^k b_{0,n} \right). \end{aligned} \quad (1.6)$$

We will present the proof of Theorem 1.1 in the next section. In Section 3, we will compute the Fourier transforms of certain minimal supported bivariate splines to illustrate the general theory, and discuss the relation to subdivision (refinement) masks for parametric spline surface rendering.

2 Proof of Theorem

The integral of the Fourier transform \hat{P}_n of a bivariate polynomial P_n restricted to the triangle is related to the hypergeometric function ${}_1F_1$, called a confluent hypergeometric function (see e.g. [9]), defined by

$${}_1F_1(\alpha_1, \beta_1; z) := 1 + \frac{\alpha_1}{\beta_1} \frac{z}{1!} + \frac{\alpha_1(\alpha_1 + 1)}{\beta_1(\beta_1 + 1)} \frac{z^2}{2!} + \frac{\alpha_1(\alpha_1 + 1)(\alpha_1 + 2)}{\beta_1(\beta_1 + 1)(\beta_1 + 2)} \frac{z^3}{3!} + \dots, \quad z \in \mathbb{C},$$

where $\alpha_1, \beta_1 \in \mathbb{C}$ with $\beta_1 \notin \{0, -1, -2, \dots\}$. Clearly,

$${}_1F_1(0, \beta_1; z) = 1, \quad {}_1F_1(\beta_1, \beta_1; z) = e^z.$$

We need the following two properties of ${}_1F_1$ for any nonnegative integers α_1, β_1 :

(i) For integers $k \geq 0, m > 0$,

$$\frac{z}{m} {}_1F_1(k+1, m+1; z) = {}_1F_1(k+1, m; z) - {}_1F_1(k, m; z), \quad z \in \mathbb{C}; \quad (2.1)$$

(ii) For integers $k \geq 0, m \geq 0, 0 \leq u \leq 1$ and $\rho \in \mathbb{C}$,

$$\int_0^{1-u} \nu^k (1-u-\nu)^m e^{\rho\nu} d\nu = \frac{k!m!}{(k+m+1)!} {}_1F_1(k+1, k+m+2; (1-u)\rho). \quad (2.2)$$

(See [9], p.1013 and p.343).

To prove Theorem 1.1, we need the following result.

Lemma 2.1 For any integer $s \geq 0$, real numbers $b_{m,j}$, and $\rho, z \in \mathbb{C}$,

$$\begin{aligned} & \sum_{m=0}^s b_{m,s-m} \frac{1}{(s+1)!} z^{s+1} {}_1F_1(m+1, s+2; \rho z) \\ &= \sum_{\ell=0}^s \frac{(-1)^\ell}{\rho^{\ell+1}} \nabla^\ell b_{s,0} \frac{z^{s-\ell}}{(s-\ell)!} e^{\rho z} - \sum_{\ell=0}^s \frac{(-1)^\ell}{\rho^{\ell+1}} \Delta^\ell b_{0,s} \frac{z^{s-\ell}}{(s-\ell)!}. \end{aligned} \quad (2.3)$$

Proof. By applying (2.1), we see that

$$\begin{aligned} \text{Left-hand side of (2.3)} &= \sum_{m=0}^s b_{m,s-m} \frac{z^s}{\rho \cdot s!} ({}_1F_1(m+1, s+1; \rho z) - {}_1F_1(m, s+1; \rho z)) \\ &= \sum_{m=0}^s b_{m,s-m} \frac{z^s}{\rho \cdot s!} {}_1F_1(m+1, s+1; \rho z) - \sum_{m=0}^s b_{m,s-m} \frac{z^s}{\rho \cdot s!} {}_1F_1(m, s+1; \rho z) \\ &= b_{s,0} \frac{z^s}{\rho \cdot s!} {}_1F_1(s+1, s+1; \rho z) + \frac{1}{\rho} \sum_{m=0}^{s-1} b_{m,s-m} \frac{z^s}{s!} {}_1F_1(m+1, s+1; \rho z) \\ &\quad - b_{0,s} \frac{z^s}{\rho \cdot s!} {}_1F_1(0, s+1; \rho z) - \sum_{m=1}^s b_{m,s-m} \frac{z^s}{\rho \cdot s!} {}_1F_1(m, s+1; \rho z) \\ &= b_{s,0} \frac{z^s}{\rho \cdot s!} e^{\rho z} + \frac{1}{\rho} \sum_{m=0}^{s-1} b_{m,s-m} \frac{z^s}{s!} {}_1F_1(m+1, s+1; \rho z) \\ &\quad - b_{0,s} \frac{z^s}{\rho \cdot s!} - \frac{1}{\rho} \sum_{m=0}^{s-1} b_{m+1,s-m-1} \frac{z^s}{s!} {}_1F_1(m+1, s+1; \rho z) \\ &= \frac{1}{\rho} b_{s,0} \frac{z^s}{s!} e^{\rho z} - \frac{1}{\rho} b_{0,s} \frac{z^s}{s!} + \frac{(-1)}{\rho} \sum_{m=0}^{s-1} \Delta b_{m,s-m} \frac{z^s}{s!} {}_1F_1(m+1, s+1; \rho z). \end{aligned}$$

Now, repeating this process, we may conclude that the left-hand side of (2.3) is given by

$$\begin{aligned} & \left(\frac{1}{\rho} b_{s,0} \frac{z^s}{s!} + \frac{(-1)}{\rho^2} \Delta b_{s-1,1} \frac{z^{s-1}}{(s-1)!} + \cdots + \frac{(-1)^\ell}{\rho^{\ell+1}} \Delta^\ell b_{s-\ell,\ell} \frac{z^{s-\ell}}{(s-\ell)!} + \cdots + \frac{(-1)^s}{\rho^{s+1}} \Delta^s b_{0,s} \right) e^{\rho z} \\ & - \left(\frac{1}{\rho} b_{0,s} \frac{z^s}{s!} + \frac{(-1)}{\rho^2} \Delta b_{0,s} \frac{z^{s-1}}{(s-1)!} + \cdots + \frac{(-1)^\ell}{\rho^{\ell+1}} \Delta^\ell b_{0,s} \frac{z^{s-\ell}}{(s-\ell)!} + \cdots + \frac{(-1)^s}{\rho^{s+1}} \Delta^s b_{0,s} \right) \\ & = \sum_{\ell=0}^s \frac{(-1)^\ell}{\rho^{\ell+1}} \nabla^\ell b_{s,0} \frac{z^{s-\ell}}{(s-\ell)!} e^{\rho z} - \sum_{\ell=0}^s \frac{(-1)^\ell}{\rho^{\ell+1}} \Delta^\ell b_{0,s} \frac{z^{s-\ell}}{(s-\ell)!}, \end{aligned}$$

where the last equality follows from the identity $\Delta^\ell b_{s-\ell,\ell} = \nabla^\ell b_{s,0}$. Hence we obtain (2.3). \blacksquare

We are now ready to prove the main result of the paper.

Proof of Theorem 1.1. By simple calculations, we have

$$\begin{aligned} \widehat{P}_n(\xi) &= \int_{\Delta A_1 A_2 A_3} e^{-i\xi \cdot \mathbf{x}} P_n(\mathbf{x}) d\mathbf{x} = 2V_T \sum_{j=0}^n \sum_{k=0}^{n-j} a_{j,k,n-j-k} \frac{n!}{j!k!(n-j-k)!} \\ &\quad \times \int_0^1 \int_0^{1-u} e^{-i\xi_1(a_3+(a_1-a_3)u+(a_2-a_3)v)} e^{-i\xi_2(b_3+(b_1-b_3)u+(b_2-b_3)v)} u^j v^k (1-u-v)^{n-j-k} dv du \\ &= 2V_T e^{-i(a_3\xi_1+b_3\xi_2)} \sum_{j=0}^n \sum_{k=0}^{n-j} a_{j,k,n-j-k} \frac{n!}{j!k!(n-j-k)!} \int_0^1 e^{\beta u} u^j \int_0^{1-u} e^{\gamma v} v^k (1-u-v)^{n-j-k} dv du \\ &= 2V_T e^{-i(a_3\xi_1+b_3\xi_2)} \sum_{j=0}^n \sum_{k=0}^{n-j} a_{j,k,n-j-k} \frac{n!}{j!(n-j+1)!} \\ &\quad \times \int_0^1 e^{\beta u} u^j (1-u)^{n-j+1} {}_1F_1(k+1, n-j+2; (1-u)\gamma) du \\ &= 2V_T n! e^{-i(a_3\xi_1+b_3\xi_2)} \sum_{j=0}^n \int_0^1 e^{\beta u} \frac{u^j}{j!} \\ &\quad \times \left\{ \sum_{k=0}^{n-j} a_{j,k,n-j-k} \frac{1}{(n-j+1)!} (1-u)^{n-j+1} {}_1F_1(k+1, n-j+2; (1-u)\gamma) \right\} du, \end{aligned}$$

where the third equality follows from (2.2). Applying Lemma 2.1 to the sum in the curly brackets of the rightmost equality with $s = n - j$, $b_{m,\ell} = a_{j,m,\ell}$ (so that

$\Delta b_{m,\ell} = \Delta_{23} a_{j,m,\ell}, \nabla b_{m,\ell} = \nabla_{23} a_{j,m,\ell}$, $z = 1 - u, \rho = \gamma$, we see that

$$\begin{aligned}
\widehat{P}_n(\xi) &= 2V_T n! e^{-i(a_3 \xi_1 + b_3 \xi_2)} \sum_{j=0}^n \int_0^1 e^{\beta u} \frac{u^j}{j!} \\
&\quad \times \left\{ \sum_{\ell=0}^{n-j} \frac{(-1)^\ell}{\gamma^{\ell+1}} \nabla_{23}^\ell a_{j,n-j,0} \frac{(1-u)^{n-j-\ell}}{(n-j-\ell)!} e^{(1-u)\gamma} - \sum_{\ell=0}^{n-j} \frac{(-1)^\ell}{\gamma^{\ell+1}} \Delta_{23}^\ell a_{j,0,n-j} \frac{(1-u)^{n-j-\ell}}{(n-j-\ell)!} \right\} du \\
&= 2V_T n! e^{-i(a_2 \xi_1 + b_2 \xi_2)} \sum_{j=0}^n \sum_{\ell=0}^{n-j} \frac{(-1)^\ell}{\gamma^{\ell+1}} \nabla_{23}^\ell a_{j,n-j,0} \frac{1}{j!(n-j-\ell)!} \int_0^1 u^j (1-u)^{n-j-\ell} e^{\alpha u} du \\
&\quad - 2V_T n! e^{-i(a_3 \xi_1 + b_3 \xi_2)} \sum_{j=0}^n \sum_{\ell=0}^{n-j} \frac{(-1)^\ell}{\gamma^{\ell+1}} \Delta_{23}^\ell a_{j,0,n-j} \frac{1}{j!(n-j-\ell)!} \int_0^1 u^j (1-u)^{n-j-\ell} e^{\beta u} du \\
&= 2V_T n! e^{-i(a_2 \xi_1 + b_2 \xi_2)} \sum_{j=0}^n \sum_{\ell=0}^{n-j} \frac{(-1)^\ell}{\gamma^{\ell+1}} \nabla_{23}^\ell a_{j,n-j,0} \frac{1}{(n-\ell+1)!} {}_1F_1(j+1, n-\ell+2; \alpha) \\
&\quad - 2V_T n! e^{-i(a_3 \xi_1 + b_3 \xi_2)} \sum_{j=0}^n \sum_{\ell=0}^{n-j} \frac{(-1)^\ell}{\gamma^{\ell+1}} \Delta_{23}^\ell a_{j,0,n-j} \frac{1}{(n-\ell+1)!} {}_1F_1(j+1, n-\ell+2; \beta) \\
&= 2V_T n! e^{-i(a_2 \xi_1 + b_2 \xi_2)} \sum_{\ell=0}^n \frac{(-1)^\ell}{\gamma^{\ell+1}} \sum_{j=0}^{n-\ell} \nabla_{23}^\ell a_{j,n-j,0} \frac{1}{(n-\ell+1)!} {}_1F_1(j+1, n-\ell+2; \alpha) \\
&\quad - 2V_T n! e^{-i(a_3 \xi_1 + b_3 \xi_2)} \sum_{\ell=0}^n \frac{(-1)^\ell}{\gamma^{\ell+1}} \sum_{j=0}^{n-\ell} \Delta_{23}^\ell a_{j,0,n-j} \frac{1}{(n-\ell+1)!} {}_1F_1(j+1, n-\ell+2; \beta).
\end{aligned}$$

Finally, applying Lemma 2.1 to the first term (the second term resp.) in the above equation with $s = n - \ell, b_{m,k} = \nabla_{23}^\ell a_{m,k+\ell,0}, z = 1, \rho = \alpha$ (with $s = n - \ell, b_{m,k} = \nabla_{23}^\ell a_{m,0,k+\ell}, z = 1, \rho = \beta$ resp.), we have

$$\begin{aligned}
\widehat{P}_n(\xi) &= 2V_T n! e^{-i(a_2 \xi_1 + b_2 \xi_2)} \sum_{\ell=0}^n \frac{(-1)^\ell}{\gamma^{\ell+1}} \\
&\quad \times \left\{ \sum_{k=0}^{n-\ell} \frac{(-1)^k}{\alpha^{k+1}} \nabla_{12}^k \nabla_{23}^\ell a_{n-\ell,\ell,0} \frac{e^\alpha}{(n-\ell-k)!} - \sum_{k=0}^{n-\ell} \frac{(-1)^k}{\alpha^{k+1}} \Delta_{12}^k \nabla_{23}^\ell a_{0,n,0} \frac{1}{(n-\ell-k)!} \right\} \\
&\quad - 2V_T n! e^{-i(a_3 \xi_1 + b_3 \xi_2)} \sum_{\ell=0}^n \frac{(-1)^\ell}{\gamma^{\ell+1}} \\
&\quad \times \left\{ \sum_{k=0}^{n-\ell} \frac{(-1)^k}{\beta^{k+1}} \nabla_{13}^k \Delta_{23}^\ell a_{n-\ell,0,\ell} \frac{e^\beta}{(n-\ell-k)!} - \sum_{k=0}^{n-\ell} \frac{(-1)^k}{\beta^{k+1}} \Delta_{13}^k \Delta_{23}^\ell a_{0,0,n} \frac{1}{(n-\ell-k)!} \right\}
\end{aligned}$$

$$\begin{aligned}
&= 2V_T \sum_{\ell=0}^n \sum_{k=0}^{n-\ell} (-1)^{k+\ell} \frac{n!}{(n-k-\ell)!} \frac{1}{\gamma^{\ell+1}} \\
&\times \left\{ \frac{1}{\alpha^{k+1}} \left(\nabla_{12}^k \nabla_{23}^\ell a_{n-\ell, \ell, 0} e^{-i(a_1 \xi_1 + b_1 \xi_2)} - \Delta_{12}^k \nabla_{23}^\ell a_{0, n, 0} e^{-i(a_2 \xi_1 + b_2 \xi_2)} \right) \right. \\
&\quad \left. - \frac{1}{\beta^{k+1}} \left(\nabla_{13}^k \Delta_{23}^\ell a_{n-\ell, 0, \ell} e^{-i(a_1 \xi_1 + b_1 \xi_2)} - \Delta_{13}^k \Delta_{23}^\ell a_{0, 0, n} e^{-i(a_3 \xi_1 + b_3 \xi_2)} \right) \right\},
\end{aligned}$$

as desired. ■

Remark 1. By applying Lemma 2.1 and following the above procedure, the main result in this paper can be extended to higher dimensions. That is, explicit formulae of the Fourier transform of Bernstein-Bézier representations of multivariate polynomials on simplexes can be derived in a similar way.

3 Application to refinable bivariate splines on triangles

Refinable spline functions are instrumental to surface subdivisions. For example, the bi-cubic B -spline is used in the Catmull-Clark scheme [1] and the three-direction box-spline B_{222} is used in the Loop scheme [10]. The simple reasons are that firstly, the refinement masks of such spline functions immediately give the so-called “local averaging rules” for the subdivision schemes; and secondly, the parametric spline representations are precisely the subdivision surfaces. While the refinement masks of the bi-cubic B -spline and the box-spline B_{222} , being defined by convolutions of the characteristic function of the unit square along the appropriate directions, are readily computable, those for others, such as basis functions with minimum and quasi-minimum supports, are not as easy to compute. Examples of the recent development in this direction are the refinable bivariate C^2 -cubic, C^2 -quartic and C^2 -quintic spline functions in [6, 7, 8], introduced for matrix-valued surface subdivisions to gain such desirable properties as surface geometric shape control parameters, smaller subdivision template size (to better address the often unavoidable extraordinary vertices), and interpolation of the position components of the (initial) control vertices. Computations of the refinement masks of these bivariate spline functions are very tedious, requiring formulating and solving large linear systems in terms of Bernstein-Bézier coefficients. For this reason, the original motivation for this research was to extend the standard Fourier approach to computing the (scalar-valued) refinement masks of refinable spline functions to computing the matrix-valued refinement masks of refinable spline function vectors.

As an application, let us consider the Fourier transform of the minimally supported (ms) and quasi-minimally supported (qms) bivariate splines in $S_{m(k)}^k(\Delta^{(i)})$ ($i = 1, 2$), where $\Delta^{(1)}$ and $\Delta^{(2)}$ denote respectively 3 and 4-direction meshes in

\mathbb{R}^2 with integer grid points, and $S_{m(k)}^k(\Delta^{(i)})$ ($i = 1, 2$) the spaces of functions in C^k whose restrictions on the triangular cells are polynomials of degree $m(k)$. Here, $m(k)$ denotes the smallest nonnegative integer such that $S_{m(k)}^k(\Delta^{(i)})$ contains at least one locally supported function f . It is well known that for $\Delta^{(1)}$ $m(2k-1) = 3k$ and $m(2k) = 3k+1$; and for $\Delta^{(2)}$ we have $m(3k-1) = 4k$, $m(3k) = 4k+1$, and $m(3k+1) = 4k+2$ (see [2, 3]).

Minimally ms and qms bivariate splines, considered in [2, 3, 4], are defined as follows. The support of a locally supported function f in a spline space is the closure of the set on which f does not vanish and is denoted by $\text{supp}(f)$. A set S is called a minimal support of a spline space if there is some f , called an ms spline, in the space with $\text{supp}(f) = S$, but there does not exist a nontrivial g in the space with $\text{supp}(g)$ properly contained in S . A function f in a spline space is called a qms spline if (i) f cannot be written as a (finite) linear combination of ms splines in the space, and (ii) for any h in the space properly contained in $\text{supp}(f)$, h is some (finite) linear combination of ms splines in the space.

For example, for the 3-direction mesh $\Delta^{(1)}$, while the spline space $S_1^0(\Delta^{(1)})$ has only one ms spline g_1^0 , which is the box spline B_{111} with direction set $\{(1, 0), (0, 1), (1, 1)\}$, the space $S_0^{-1}(\Delta^{(1)})$ has two ms splines g_2^0 and g_3^0 which are the characteristic functions of χ_A and χ_B , where A is the triangle bounded by the 3 lines: $x = 0$, $y = 1$, and $y = x$, and B the triangle bounded by the 3 lines: $x = 1$, $y = 0$ and $y = x$.

On the other hand, for the 4-direction mesh $\Delta^{(2)}$, the space $S_2^1(\Delta^{(2)})$ has only one ms spline $f_1^0 = B_{1111}$, the box spline with direction set $\{(1, 0), (0, 1), (1, 1), (1, -1)\}$, but there are two ms splines in the space $S_1^0(\Delta^{(2)})$ by f_2^0 and f_3^0 , where f_2^0 is the Courant hat function having a diamond support with the vertices $(1, 0)$, $(0, 1)$, $(-1, 0)$, and $(0, -1)$, and having value of 1 at the center $(0, 0)$ of the support, and f_3^0 the other Courant hat function supported on the unit square $[0, 1]^2$ with value 1 at its center $(1/2, 1/2)$. Furthermore, there are two ms splines f_4^0 and f_5^0 and one qms spline f_6^0 in $S_4^2(\Delta^{(2)})$ (see [2, 3, 4]).

In the following, let us consider the k -fold 2-dimensional convolutions: $g_1^k = \underbrace{g_1^0 * g_1^0 * \cdots * g_1^0}_{k+1}$ and $f_1^k = \underbrace{f_1^0 * f_1^0 * \cdots * f_1^0}_{k+1}$ of g_1^0 and f_1^0 , respectively. Now, the spline function vectors of interest are:

$$G^k \equiv \begin{bmatrix} g_2^k \\ g_3^k \end{bmatrix} := g_1^{k-1} * \begin{bmatrix} g_2^0 \\ g_3^0 \end{bmatrix}, \quad F^k \equiv \begin{bmatrix} f_2^k \\ f_3^k \end{bmatrix} := f_1^{k-1} * \begin{bmatrix} f_2^0 \\ f_3^0 \end{bmatrix}, \quad (3.1)$$

where $k \geq 1$ and $f * \begin{bmatrix} g \\ h \end{bmatrix} := \begin{bmatrix} f * g \\ f * h \end{bmatrix}$.

We remark that among the functions g_i^k, f_i^k , $i = 1, 2, 3$, only g_1^k and f_1^k are box splines, while all of them are the unique ms and qms splines in the corresponding

spline spaces (where the notion of uniqueness is according to the statement of Theorem 3.2 in [2]).

The Fourier transforms of the “initial” ms splines $g_i^0, f_i^0, i = 2, 3$, can be evaluated by using the formula provided in this paper, and the Fourier transforms of the other splines are given by the corresponding products with those of g_1^k or f_1^k .

Finally, the refinement masks of G^k, F^k can be easily computed by making use of (3.1) from the refinements of the “initial” G^0 and F^0 .

Example 1. The Fourier transform of G^k is given by $\widehat{G^k}(\xi_1, \xi_2) = [\widehat{g_2^k}, \widehat{g_3^k}]^T(\xi_1, \xi_2)$, where $\widehat{g_2^k} = \widehat{g_2^0}(\widehat{g_1^0})^k$ and $\widehat{g_3^k} = \widehat{g_3^0}(\widehat{g_1^0})^k$. Here, we have

$$\begin{aligned}\widehat{g_1^0}(\xi_1, \xi_2) &= \widehat{B_{111}}(\xi_1, \xi_2) = \frac{1 - e^{-i\xi_1}}{i\xi_1} \frac{1 - e^{-i\xi_2}}{i\xi_2} \frac{1 - e^{i(\xi_1 + \xi_2)}}{i(\xi_1 + \xi_2)}, \\ \widehat{g_2^0}(\xi_1, \xi_2) &= \frac{1 - e^{-i(\xi_1 + \xi_2)}}{\xi_1(\xi_1 + \xi_2)} - \frac{1 - e^{-i\xi_2}}{\xi_1\xi_2},\end{aligned}\tag{3.2}$$

and

$$\widehat{g_3^0}(\xi_1, \xi_2) = \frac{1 - e^{-i(\xi_1 + \xi_2)}}{\xi_2(\xi_1 + \xi_2)} - \frac{1 - e^{-i\xi_1}}{\xi_1\xi_2}.\tag{3.3}$$

■

Next, let us compute the Fourier transform of F^k . For this purpose, recall that

$$(\phi(A \cdot -\mathbf{k}))^\wedge(\xi) = |\det(A)|^{-1} e^{-i\xi^T \cdot (A^{-1}\mathbf{k})} \widehat{\phi}((A^{-1})^T \xi),\tag{3.4}$$

for any invertible matrix A of dimension s .

Example 2. Let f_2^0, f_3^0 be the two Courant hat functions in $S_1^0(\Delta^{(2)})$ as introduced previously. To compute the Fourier transform of f_2^0 , we use the x - y axes to partition its support into four triangles: $\Delta_1, \Delta_2, \Delta_3$, and Δ_4 in the first, second, third, and fourth quadrants, respectively. Then f_2^0 can be written as the sum of four functions: ϕ_1, ϕ_2, ϕ_3 , and ϕ_4 , with supports given by $\Delta_1, \Delta_2, \Delta_3$, and Δ_4 , respectively.

By the formula (1.3), the Fourier transform of ϕ_1 is given by

$$\widehat{\phi_1}(\xi_1, \xi_2) = -\frac{1}{\xi_1\xi_2} + i\frac{1 - e^{-i\xi_1}}{\xi_1^2(\xi_1 - \xi_2)} + i\frac{1 - e^{-i\xi_2}}{\xi_2^2(\xi_2 - \xi_1)}.\tag{3.5}$$

Since $\phi_2(x, y) = \phi_1(-x, y)$, it follows from (3.5) and (3.4) that

$$\widehat{\phi_2}(\xi_1, \xi_2) = \widehat{\phi_1}(-\xi_1, \xi_2) = \frac{1}{\xi_1\xi_2} - i\frac{1 - e^{i\xi_1}}{\xi_1^2(\xi_1 + \xi_2)} + i\frac{1 - e^{-i\xi_2}}{\xi_2^2(\xi_1 + \xi_2)}.$$

Similarly, since $\phi_3(x, y) = \phi_1(-x, -y)$ and $\phi_4(x, y) = \phi_1(x, -y)$, we have

$$\begin{aligned}\widehat{\phi}_3(\xi_1, \xi_2) &= -\frac{1}{\xi_1 \xi_2} + i \frac{1 - e^{i\xi_1}}{\xi_1^2(\xi_2 - \xi_1)} + i \frac{1 - e^{i\xi_2}}{\xi_2^2(\xi_1 - \xi_2)}; \\ \widehat{\phi}_4(\xi_1, \xi_2) &= \frac{1}{\xi_1 \xi_2} + i \frac{1 - e^{-i\xi_1}}{\xi_1^2(\xi_1 + \xi_2)} - i \frac{1 - e^{i\xi_2}}{\xi_2^2(\xi_1 + \xi_2)}.\end{aligned}$$

Consequently, we arrive at

$$\widehat{f}_2^0(\xi_1, \xi_2) = \sum_{j=1}^4 \widehat{\phi}_j(\xi_1, \xi_2) = 2i \frac{e^{i\xi_2} - e^{-i\xi_2}}{\xi_2(\xi_1 + \xi_2)(\xi_2 - \xi_1)} + 2i \frac{e^{i\xi_1} - e^{-i\xi_1}}{\xi_1(\xi_1 + \xi_2)(\xi_1 - \xi_2)}. \quad (3.6)$$

Next, observe that the linear transformation $B \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, with

$$B = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix},$$

maps the $\text{supp}(f_3^0)$ to $\text{supp}(f_2^0)$, with the vertices $(0, 0)$, $(1, 0)$, $(1, 1)$, and $(0, 1)$ of $\text{supp}(f_3^0)$ corresponding to the vertices $(-1, 0)$, $(0, 1)$, $(1, 0)$, and $(0, -1)$ of $\text{supp}(f_2^0)$, respectively. Hence, we may write

$$f_3^0(x, y) = f_2^0\left(B \begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} 1 \\ 0 \end{bmatrix}\right),$$

and apply (3.4) to obtain

$$\begin{aligned}\widehat{f}_3^0(\xi_1, \xi_2) &= \frac{1}{2} e^{-i\xi \cdot \frac{1}{2} B[1, 0]^T} \widehat{f}_2^0\left(\frac{1}{2} B\xi\right) = \frac{1}{2} e^{-i\frac{\xi_1 + \xi_2}{2}} \widehat{f}_2^0\left(\frac{\xi_1 + \xi_2}{2}, \frac{\xi_1 - \xi_2}{2}\right) \\ &= \frac{2i}{\xi_1 \xi_2} \left(\frac{e^{-i\xi_2} - e^{-i\xi_1}}{\xi_2 - \xi_1} + \frac{1 - e^{-i(\xi_1 + \xi_2)}}{\xi_1 + \xi_2} \right).\end{aligned} \quad (3.7)$$

■

Finally, to compute the refinement masks of G^k and F^k in (3.1), we observe that the convolution of two (finitely) refinable functions remains to be (finitely) refinable, and that G^k is the convolution of G^0 with the refinable box spline g_1^k , and F^k the convolution of F^0 with the refinable box spline f_1^k . Hence, we only need to compute the refinement masks of the initial ms splines. Precisely, from \widehat{g}_2^0 and \widehat{g}_3^0 given by (3.2) and (3.3), we have

$$\widehat{G}^0(2\xi_1, 2\xi_2) = Q_0(\xi_1, \xi_2) \widehat{G}^0(\xi_1, \xi_2)$$

where

$$Q_0(\xi_1, \xi_2) = \begin{bmatrix} 1 + z_2 + z_1 z_2 & z_2 \\ z_1 & 1 + z_1 + z_1 z_2 \end{bmatrix}, \quad z_1 = e^{-i\xi_1}, \quad z_2 = e^{-i\xi_2};$$

and from \hat{f}_2^0 and \hat{f}_3^0 given by (3.6) and (3.7), we have

$$\hat{F}^0(2\xi_1, 2\xi_2) = R_0(\xi_1, \xi_2)\hat{F}^0(\xi_1, \xi_2)$$

where

$$R_0(\xi_1, \xi_2) = \frac{1}{4} \begin{bmatrix} 1 + \frac{1}{2}(z_1 + \frac{1}{z_2})(z_1 + z_2) & \frac{1}{2}(1 + \frac{1}{z_1})(1 + \frac{1}{z_2}) \\ z_1 z_2 & \frac{1}{2}(1 + z_1)(1 + z_2) \end{bmatrix}, \quad z_1 = e^{-i\xi_1}, z_2 = e^{-i\xi_2}.$$

The interested reader is referred to [5] for computing the refinement mask for $F^0(\mathbf{x})$ by calculating the BB coefficients of $F^0(B^{-1}\mathbf{x})$ directly.

Therefore, we conclude, from the definitions in (3.1), that

$$\hat{G}^k(2\xi_1, 2\xi_2) = Q_k(\xi_1, \xi_2)\hat{G}^k(\xi_1, \xi_2), \quad \hat{F}^k(2\xi_1, 2\xi_2) = R_k(\xi_1, \xi_2)\hat{F}^k(\xi_1, \xi_2)$$

with

$$Q_k(\xi_1, \xi_2) = (q(\xi_1, \xi_2))^k Q_0(\xi_1, \xi_2), \quad R_k(\xi_1, \xi_2) = (r(\xi_1, \xi_2))^k R_0(\xi_1, \xi_2) \quad (3.8)$$

where $q(\xi_1, \xi_2) = \frac{1}{8}(1 + e^{-i\xi_1})(1 + e^{-i\xi_2})(1 + e^{-i(\xi_1+\xi_2)})$ is the mask (or two-scale symbol) of g_1^0 , and $r(\xi_1, \xi_2) = \frac{1}{16}(1 + e^{-i\xi_1})(1 + e^{-i\xi_2})(1 + e^{-i(\xi_1+\xi_2)})(1 + e^{-i(\xi_1-\xi_2)})$ the mask (or two-scale symbol) of f_1^0 .

That is, we have the following result.

Theorem 3.1 *The vector-valued functions G^k and F^k are finitely refinable with refinement masks given by (3.8).*

Similarly,

$$f_1^{k-1} * \begin{bmatrix} f_4^0 \\ f_5^0 \\ f_6^0 \end{bmatrix}$$

is also finitely refinable, with refinement mask given by

$$(r(\xi_1, \xi_2))^k S_0(\xi_1, \xi_2),$$

where S_0 is the refinement mask of $\begin{bmatrix} f_4^0 \\ f_5^0 \\ f_6^0 \end{bmatrix}$. Computation of S_0 as well as the refinement masks of other initial ms and qms bivariate splines in general is usually nontrivial and requires further investigation.

References

- [1] E. Catmull and J. Clark, Recursively generated B-splines surfaces on arbitrary topological meshes, *Comput. Aided Design* **10** (1978), 350–355.
- [2] C.K. Chui, *Multivariate Splines*, NSF-CBMS Series #54, SIAM Publ., Philadelphia, 1988.
- [3] C.K. Chui and T.X. He, Computation of minimally and quasi-minimally supported bivariate splines, *J. Comput. Math* **8** (1990), 108–117.
- [4] C.K. Chui and T.X. He, On minimal and quasi-minimal supported bivariate splines, *J. Approx. Theory* **52** (1998), 217–238.
- [5] C.K. Chui and Q.T. Jiang, Multivariate balanced vector-valued refinable functions, In: *Modern Development in Multivariate Approximation*, ISNM 145, V.W. Haussmann, K. Jetter, M. Reimer, and J. Stöckler (eds.), Birkhäuser Verlag, Basel, 2003, pp. 71–102.
- [6] C.K. Chui and Q.T. Jiang, Surface subdivision schemes generated by refinable bivariate spline function vectors, *Appl. Comput. Harmon. Anal.* **15** (2003), 147–162.
- [7] C.K. Chui and Q.T. Jiang, Refinable bivariate C^2 -splines for multi-level data representation and surface display, *Math Comp.* **74** (2005), 1369–1390.
- [8] C.K. Chui and Q.T. Jiang, Refinable bivariate quartic and quintic C^2 -splines for quadrilateral subdivisions, *J. Comp. Appl. Math.*, to appear. Available: www.cs.ums1.edu/~jiang
- [9] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series, and Products*, 6th edition, Academic Press, New York, 2000.
- [10] C. Loop, Smooth subdivision surfaces based on triangles, Master’s Thesis, University of Utah, Department of Mathematics, Salt Lake City, 1987.